

## SYNCHROTRON-BASED HIGH RESOLUTION FAR INFRARED SPECTROSCOPY OF BENZALDEHYDE

YUE LIANG, ZENGKUI LIU, JIARUI MA, YICHI ZHANG, HAIHUA ZHOU, ZIQIU CHEN, *College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou, Gansu, China*; CSABA FÁBRI, *Laboratory of Molecular Structure and Dynamics, Eötvös University, Budapest, Hungary*; BRANT E. BILLINGHURST, JIANBAO ZHAO, *EFD, Canadian Light Source Inc., Saskatoon, Saskatchewan, Canada*.

The rotationally-resolved vibrational spectra of benzaldehyde have been recorded in the far infrared region at room temperature using the Bruker IFS125 Fourier Transform spectrometer at the Canadian Light Source with a resolution of  $0.000959\text{ cm}^{-1}$ . The lowest frequency vibrational bands collected correspond to the fundamental and hotbands of the -CHO torsion at  $110\text{ cm}^{-1}$ . The assignment and analysis of the dense spectral features in this region will be detailed. In addition, the newly obtained band origin of the -CHO torsional fundamental will be compared to *ab initio* results in relation to the discrepancy<sup>a</sup> between theoretical and experimental values of the potential barrier to internal rotation in this molecule.

---

<sup>a</sup>L.D. Speakman, B.N. Papas, H. Lee Woodcock, H.F. Schaefer III, *J. Chem. Phys.*, **2004**, 120, 4247-4250; I.A. Godunov, V.A. Bataev, A. V. Abramnikov, V.I. Pupyshev, *J. Phys. Chem.A*, **2014**, 118, 10159-10165, and references therein